

RKKY interaction in framework of $T = 0$ Green function method

Todor M. Mishonov, Tihomir I. Valchev, Liliya A. Atanasova, and Peter A. Ivanov

*Department of Theoretical Physics, Faculty of Physics,
Sofia University St. Kliment Ohridski, Bulgaria.*

A methodical derivation of RKKY interaction in framework of $T = 0$ Green function method is given in great detail. The article is complimentary to standard textbooks on the physics of magnetism and condensed matter physics. It is shown that the methods of statistical mechanics gives a standard and probably simplest derivation of the exchange interaction. A parallel with theory of plasma waves demonstrates the relation between the Fourier transformation of polarization operator of degenerate electron gas at zero frequency and the space dependence of the indirect electron exchange due to itinerant electrons.

PACS numbers: 71.45.Gm, 75.10.Lp, 75.30.Et

The RKKY interaction is a basic ingredient in the physics of magnetism, see for example the textbook by Mattis [1]. The purpose of the present paper is purely methodical: to derive RKKY result using the Green function technique for $T = 0$. The diagram formalism is described in the monograph by Lifshitz and Pitaevski [2] or the monograph by Abrikosov, Gor'kov and Dzyaloshinski [3]. The s-d exchange interaction

$$\hat{H}_{s-d} = -J_{s-d} (\hat{\mathbf{s}}(\mathbf{R}_1) \cdot \mathbf{S}_1 + \hat{\mathbf{s}}(\mathbf{R}_2) \cdot \mathbf{S}_2) \quad (1)$$

is treated as a small perturbation. Where $\mathbf{R}_1$ and $\mathbf{R}_2$ are the space-vectors of the impurities, $\mathbf{S}_1$ and $\mathbf{S}_2$ are their spins, and $\hat{\mathbf{s}}(\mathbf{R}_1)$ and $\hat{\mathbf{s}}(\mathbf{R}_2)$ are electron spin operators in the corresponding space points. Using the simple matrix equation

$$\text{Tr}(\sigma \cdot \mathbf{S}_1 \sigma \cdot \mathbf{S}_2) = 2\mathbf{S}_1 \cdot \mathbf{S}_2 \quad (2)$$

and the general rules for single loop approximation we obtain the space dependence of the indirect exchange

$$\hat{H}_{\text{RKKY}} = -J(\mathbf{R})\mathbf{S}_1 \cdot \mathbf{S}_2, \quad \mathbf{R} = \mathbf{R}_2 - \mathbf{R}_1, \quad R = |\mathbf{R}|, \quad (3)$$

where

$$J(R) = \left(\frac{J_{s-d}}{2} \right)^2 \left(\frac{a_0}{2\pi} \right)^6 \int_{E(k) < \mu} d^3k \int_{E(k') > \mu} d^3k' \frac{2 \cos((\mathbf{k} - \mathbf{k}') \cdot \mathbf{R})}{\frac{\hbar^2}{2m^*}(\mathbf{k}'^2 - \mathbf{k}^2 + i2m^*/\hbar\tau)}, \quad (4)$$

where $\mu = \frac{k_F^2 \hbar^2}{2m^*}$ is the Fermi energy and $\hbar k_F$ is the Fermi momentum. This integral is actually the Fourier transformation of the polarization operator of 3 dimensional electron gas for zero frequency. For a complementary Green function derivation of the polarizability of degenerate electron gas see the recent preprint [4].

For simplification of further notation we introduce the constant which parameterize the electron scattering rate $q^2 = 2m^*/\hbar\tau$. Using the elementary relation

$$\cos((\mathbf{k} - \mathbf{k}') \cdot \mathbf{R}) = \cos(\mathbf{k} \cdot \mathbf{R}) \cos(\mathbf{k}' \cdot \mathbf{R}) + \sin(\mathbf{k} \cdot \mathbf{R}) \sin(\mathbf{k}' \cdot \mathbf{R})$$

the integral above can be presented as

$$\int_{E(k) < \mu} d^3k \int_{E(k') > \mu} d^3k' \frac{\cos((\mathbf{k} - \mathbf{k}') \cdot \mathbf{R})}{\mathbf{k}'^2 - \mathbf{k}^2 + iq^2} = I_1 + I_2. \quad (5)$$

In order to solve these integrals we introduce spherical coordinates in momentum spaces

$$\mathbf{k} = k(\cos \varphi \sin \theta, \sin \varphi \sin \theta, \cos \theta), \quad \mathbf{R} = R(0, 0, 1),$$

$$d^3k = k^2 \sin \theta dk d\theta d\varphi, \quad \mathbf{k} \cdot \mathbf{R} = k R \cos \theta.$$

One can easily check that due to the symmetry of integrant the integration on k' can be expanded to $k' < k_F$. Then

$$I_1 = \int_0^{2\pi} \int_0^\pi \int_0^{k_F} d\varphi d\theta dk k^2 \sin \theta \int_0^{2\pi} \int_0^\pi \int_0^\infty d\varphi' d\theta' dk' k'^2 \sin \theta' \quad (6)$$

$$\times \frac{\sin(\mathbf{k} \cdot \mathbf{R}) \sin(\mathbf{k}' \cdot \mathbf{R})}{\mathbf{k}'^2 - \mathbf{k}^2 + iq^2} \quad (7)$$

$$= (2\pi)^2 \int_0^\pi \int_0^{k_F} d\theta dk k^2 \sin \theta \sin(k R \cos \theta) \quad (8)$$

$$\times \int_0^\infty dk' \frac{k'^2}{\mathbf{k}'^2 - \mathbf{k}^2 + iq^2} \int_0^\pi d\theta' \sin \theta' \sin(k' R \cos \theta). \quad (9)$$

And using

$$\int_0^\pi d\theta \sin \theta \sin(k R \cos \theta) = \frac{1}{k R} \cos(k R \cos \theta) \Big|_\pi^0 = 0, \quad (10)$$

we find that

$$I_1 = 0. \quad (11)$$

Analogously

$$I_2 = \int_0^{2\pi} \int_0^\pi \int_0^{k_F} d\varphi d\theta dk k^2 \sin \theta \int_0^{2\pi} \int_0^\pi \int_0^\infty d\varphi' d\theta' dk' k'^2 \sin \theta' \quad (12)$$

$$\times \frac{\cos(\mathbf{k} \cdot \mathbf{R}) \cos(\mathbf{k}' \cdot \mathbf{R})}{\mathbf{k}'^2 - \mathbf{k}^2 + iq^2} \quad (13)$$

$$= (2\pi)^2 \int_0^\pi \int_0^{k_F} d\theta dk k^2 \sin \theta \cos(k R \cos \theta) \quad (14)$$

$$\times \int_0^\infty dk' \frac{k'^2}{\mathbf{k}'^2 - \mathbf{k}^2 + iq^2} \int_0^\pi d\theta' \sin \theta' \cos(k' R \cos \theta) \quad (15)$$

and

$$\int_0^\pi d\theta \sin \theta \cos(k R \cos \theta) = \frac{1}{k R} \sin(k R \cos \theta) \Big|_\pi^0 = \frac{2 \sin(k R)}{k R} \quad (16)$$

we get

$$I_2 = \left(\frac{4\pi}{R} \right)^2 \int_0^{k_F} dk k \sin(k R) \int_0^\infty dk' \frac{k' \sin(k' R)}{\mathbf{k}'^2 - \mathbf{k}^2 + iq^2}. \quad (17)$$

In order to calculate the last integral we will apply the residuum theorem for the function

$$f(z) = \frac{z \exp(izR)}{z^2 - k^2 + iq^2}, \quad (18)$$

i.e.

$$\oint_\Gamma f(z) dz = 2\pi i \sum_n \text{Res} f(z_n), \quad 1/f(z_n) = 0, \quad (19)$$

we suppose that function f has simple poles. The contour contains the real axe and infinite arch in upper semiplane; $k' = \Re z$. The arch gives zero part when it radius goes to infinity and the real axe integral just presented by the residuum

$$\int_{-\infty}^\infty f(z) dz = 2\pi i \sum_n \text{Res} f(z_n). \quad (20)$$

The poles of $f(z)$ obeys the equation

$$z^2 = k^2 - iq^2,$$

which has the solutions

$$z_{1,2} = \sqrt[4]{k^4 + q^4} e^{i(\alpha + n\pi)}, \quad 2\alpha = -\arctan \frac{q^2}{k^2} \quad n = 0, 1. \quad (21)$$

Only $z = z_2$ belongs to upper semiplane and its residuum is

$$\text{Res}f(z = z_2) = \lim_{z \rightarrow z_2} (z - z_2)f(z) = \frac{z_2 e^{iz_2 R}}{z_2 - z_1} = \frac{e^{iz_2 R}}{2}. \quad (22)$$

In weak scattering limit

$$q \ll k_F, \quad |2\alpha| = \frac{q^2}{k^2} \ll 1, \quad z_2 \approx -k(1 + i\frac{q^2}{2k^2})$$

and

$$\text{Res}f(z_2) \approx \frac{1}{2}e^{-ikR}e^{-q^2 R/2k}. \quad (23)$$

Then

$$\int_{-\infty}^{\infty} f(z)dz = i\pi e^{-ikR}e^{-q^2 R/2k} \quad (24)$$

and

$$\int_0^{\infty} dk' \frac{k' \sin(k' R)}{\mathbf{k}'^2 - \mathbf{k}^2 + iq^2} = \frac{1}{2} \Im \left(\int_{-\infty}^{\infty} f(z)dz \right) = \frac{\pi}{2} \cos(k R) e^{-q^2 R/2k}. \quad (25)$$

In the same weak scattering limit

$$\frac{q^2}{2k} \approx \frac{q^2}{2k_F} = \frac{1}{\lambda}$$

where $\lambda = \hbar k_F \tau / m^*$ is the electron mean free path, we have

$$I_2 = \left(\frac{4\pi}{R} \right)^2 \frac{\pi}{2} e^{-R/\lambda} \int_0^{k_F} dk k \sin(k R) \cos(k R). \quad (26)$$

The elementary integration gives

$$\int_0^{k_F} dk k \sin(k R) \cos(k R) = \frac{1}{2} \int_0^{k_F} dk k \sin(2k R) \quad (27)$$

$$= -\frac{k_F \cos(2k_F R)}{4R} + \frac{1}{4R} \int_0^{k_F} dk \cos(2k R) \quad (28)$$

$$= \frac{\sin(2k_F R)}{8R^2} - \frac{k_F \cos(2k_F R)}{4R}. \quad (29)$$

And for the second integral we obtain

$$I_2 = \left(\frac{4\pi}{R} \right)^2 \frac{\pi}{2} e^{-R/\lambda} \frac{\sin(2k_F R) - 2k_F R \cos(2k_F R)}{8R^2}. \quad (30)$$

Finally for the exchange integral we arrive at

$$J(R) = (J_{s-d})^2 \left(\frac{a_0}{2\pi} \right)^6 \frac{\pi m^*}{2\hbar^2} \left(\frac{4\pi}{R} \right)^2 e^{-R/\lambda} \quad (31)$$

$$\times \frac{\sin(2k_F R) - 2k_F R \cos(2k_F R)}{8R^2} \quad (32)$$

$$= (J_{s-d})^2 \left(\frac{a_0 k_F}{2} \right)^6 \frac{8}{\pi^3 \mu} e^{-R/\lambda} \quad (33)$$

$$\times \frac{\sin(2k_F R) - 2k_F R \cos(2k_F R)}{(2k_F R)^4} \quad (34)$$

Comparing with other explanations we come to the conclusion that the method of $T = 0$ Green functions give a standard and probably simplest explanation of the RKKY interaction in metallic alloys. A derivation directly based on the methods of statistical mechanics. If we used temperature Green functions applied to $T \rightarrow 0$ we would arrive at the the same residuum.

-
- [1] D.C. Mattis, *The Theory of Magnetism*, (Harper & Row, New York, 1965), Chap. 7.
 - [2] E.M. Lifshitz and L.P. Pitaevskii, *Course on Theoretical Physics, Vol. 9, Statistical Physics, Part 2*, (Pergamon Press, Oxford, 1981), Chap. 2, Sec. 13
 - [3] A.A. Abrikosov, L.P. Gor'kov and I.E. Dzyaloshinski, *Methods of Quantum Field Theory in Statistical Physics*, (Dover, New York, 1963).
 - [4] T.M. Mishonov, L.A. Atanasova, P.A. Ivanov, T.I. Valchev *Plasma spectrum of degenerated electron gas. $T = 0$ Green function method*, cond-mat.